

# Low-energy Effective World-sheet Theory of a Non-Abelian Vortex in High Density QCD Revisited: A Regular Gauge Construction

Chandrasekhar Chatterjee<sup>1,\*</sup> and Muneto Nitta<sup>1,†</sup>

<sup>1</sup>*Department of Physics, and Research and Education Center for Natural Sciences,  
Keio University, Hiyoshi 4-1-1, Yokohama, Kanagawa 223-8521, Japan*

Color symmetry is spontaneously broken in quark matter at high density as a consequence of di-quark condensations with exhibiting color superconductivity. Non-Abelian vortices or color magnetic flux tubes stably exist in the color-flavor locked phase at asymptotically high density. The effective world-sheet theory of a single non-Abelian vortex was previously calculated in the singular gauge to obtain the  $\mathbb{CP}^2$  model [1, 2]. Here, we reconstruct the effective theory in a regular gauge without taking a singular gauge, confirming the previous results in the singular gauge.

## I. INTRODUCTION

Quark matter at high temperature and/or high density is one of important subjects both in theoretical and experimental point of view. At high density, quark matter are expected to condensate by constituting Cooper pairs. Then, color symmetry is spontaneously broken, with exhibiting color superconductivity. [3, 4], see Refs. [5, 6] as a review. The two flavor pairing may occurs in two-SC phase in which up and down quarks participate in condensations at intermediate density. At asymptotically high densities, if we can neglect the strange quark mass, the system possesses an  $SU(3)$  global flavor symmetry. In that region, it may be possible to have a three flavor pairing state, which is known as the “color-flavor locked (CFL)” phase, in which up, down and strange quarks participate in condensations. The Ginzburg-Landau (GL) free energy [7–9] shows that in the CFL ground state the baryon number  $U(1)_B$ , color  $SU(3)_C$  and flavor  $SU(3)_F$  symmetries are spontaneously broken down to the diagonal subgroup  $SU(3)_{C+F}$ . In particular,  $U(1)_B$  and color symmetry breakings lead to superfluidity and color superconductivity, respectively. Therefore, when the CFL medium rotates,  $U(1)_B$  superfluid vortices with the quantized circulations are created along the rotation axis [7, 8, 10] as the case of helium superfluids and ultra-cold atomic gases. Compared to the quantized unit circulation of  $U(1)_B$  superfluid vortices, vortices with smaller circulations ( $1/3$  quantized circulations) exist, which also carry color magnetic fluxes. They are non-Abelian vortices or color magnetic flux tubes [11–14], see Ref. [15] for a review. It was conjectured that one  $U(1)_B$  superfluid vortex is energetically split into a set of three color flux tubes with total color cancelled out [12], which has been recently confirmed numerically [16].

One non-Abelian vortex breaks the color-flavor symmetry  $SU(3)_{C+F}$  further into its subgroup in the vicinity of the core, generating Nambu-Goldstone modes (or collective coordinates) which parametrize a complex projective space  $\mathbb{CP}^2 \simeq SU(3)_{C+F}/[SU(2) \times U(1)]$ . These  $\mathbb{CP}^2$  modes are localized around the vortex core and propagate along the vortex line as gapless excitations [1, 2]. A lot of rich physics were obtained from these  $\mathbb{CP}^2$  modes. When the coupling of the  $\mathbb{CP}^2$  target space to electro-magnetic fields is introduced [17], it implies that a vortex lattice system behaves as a polarizer [18]. It also shows the Aharonov-Bohm scattering of charged particles such as electrons and muons [19]. The quantum mechanically induced gap shows the confinement of monopoles in the CFL phase, that is, quark condensations lead monopole confinement [20, 21], which gives an evidence of hadron-quark duality to the confinement phase in which quark confinement is expected to occur due to monopole condensations. Vortices interact with gluons by a topological interaction [22], implying that in the system of multiple vortices such as a vortex lattice,  $\mathbb{CP}^2$  modes in individual vortices are aligned by the interaction, exhibiting color ferromagnetism [23]. Even though such discoveries of rich physics, there remains one technical problem in the derivation of the effective Lagrangian in Refs. [1, 2]; in these references a singular gauge was taken to construct the effective  $\mathbb{CP}^2$  model. It is well known that in general one needs a careful treatment in the singular gauge. For instance, in the Abelian-Higgs model relevant for conventional metallic superconductors, a magnetic flux of a vortex is unphysically removed by taking the singular gauge. Therefore, one needs carefully to check whether the results in Refs. [1, 2] for the effective action is correct and whether no additional term exists or not.

In this paper, we construct the effective Lagrangian of a single non-Abelian vortex in a regular gauge without taking a singular gauge, and confirm that the result of the effective Lagrangian calculated in the singular gauge in Refs. [1, 2] is correct and no further term exists. To do this, we generalize the Ansatz for the gauge field used in the singular gauge, because it does not solve the Gauss’s law constraint in the regular gauge; We introduce two different profile

\* chandra@phys-h.keio.ac.jp

† nitta@phys-h.keio.ac.jp

functions that depend on both the radial coordinate  $r$  and the azimuthal angle  $\theta$ , in contrast to the singular gauge for which the profile function depends only on  $r$ . The profile functions of the zero modes are expanded in terms of partial waves, and we check the asymptotic behaviors of all the partial wave modes. By inserting the solutions of the partial wave modes into the original GL action, we derive the effective action of the vortex as the  $\mathbb{C}P^2$  action living on the vortex world-sheet. The mode previously found in the singular gauge comes out as a normalizable mode among the partial wave modes discussed in this paper. By showing the rests of partial modes are all non-normalizable, we prove the previous result on the effective theory on the vortex is correct.

This paper is organized as follows. In Sec. II, we review the GL effective theory, a non-Abelian vortex solution and its properties. In Sec. III we construct the effective theory of a single vortex in a regular gauge. Sec. IV is devoted to a summary and discussion.

## II. THE GINZBURG-LANDAU DESCRIPTION OF DENSE QCD AND A NON-ABELIAN VORTEX

### A. Ginzburg-Landau effective theory

We start with time dependent GL Lagrangian for the CFL order parameters  $\Phi_L$  and  $\Phi_R$  which are defined as di-quark condensates

$$\Phi_{La}^A \sim \epsilon_{abc} \epsilon^{ABC} q_{Lb}^B \mathcal{C} q_{Lc}^C, \quad \Phi_{Ra}^A \sim \epsilon_{abc} \epsilon^{ABC} q_{Rb}^B \mathcal{C} q_{Rc}^C, \quad (1)$$

where  $q_{L/R}$  stand for left and right handed quarks with  $a, b, c$  as fundamental color ( $SU(3)_C$ ),  $A, B, C$  as fundamental flavor ( $SU(3)_{L/R}$ ) indices and  $\mathcal{C}$  is the charge conjugation operator. Since at a high density region a perturbative calculation shows mixing terms between  $\Phi_L$  and  $\Phi_R$  are negligible, we simply assume  $\Phi_L = -\Phi_R = \Phi$  and fix their relative phase to unity. The Lagrangian has been obtained as a low-energy effective theory of the high density QCD in the CFL phase [7–9, 24]<sup>1</sup>

$$\mathcal{L}_{GL} = \text{Tr} \left[ -\epsilon_3 F_{0i} F^{0i} - \frac{1}{2\lambda_3} F_{ij} F_{ij} + K_0 \nabla_0 \Phi^\dagger \nabla_0 \Phi - K_3 \nabla_i \Phi^\dagger \nabla_i \Phi - V(\Phi) \right],$$

$$V(\phi) = -m^2 \Phi^\dagger \Phi + \beta \left[ (\text{Tr}[\Phi^\dagger \Phi])^2 + \text{Tr} \{ (\Phi^\dagger \Phi)^2 \} \right] + \frac{3m^4}{16\beta}, \quad (2)$$

where  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ig_s [A_\mu, A_\nu]$ ,  $\nabla_\mu = \partial_\mu - ig_s A_\mu$ ,  $\mu = 0, 1, 2, 3$  is the space-time index,  $\{i, j\} = \{1, 2, 3\}$  are spatial indices,  $\lambda_3$  is a magnetic permeability and  $\epsilon_3$  is a dielectric constant for gluons. Here we ignore the strange quark mass. Then the static GL free energy functional can be written as

$$-\mathcal{L}_{GL} = \text{Tr} \left[ \frac{1}{2\lambda_3} F_{ij} F^{ij} + K_3 \nabla_i \Phi^\dagger \nabla_i \Phi + V(\Phi) \right], \quad V(\phi) = -m^2 \Phi^\dagger \Phi + \beta \left[ (\text{Tr}[\Phi^\dagger \Phi])^2 + \text{Tr} \{ (\Phi^\dagger \Phi)^2 \} \right] + \frac{3m^4}{16\beta}. \quad (3)$$

The coefficients in the expression above may be calculated directly from the QCD Lagrangian using perturbative techniques. We quote here the standard results obtained in the literature [7–9] through perturbative calculations in QCD as  $\beta = \frac{7\zeta(3)}{8(\pi T_c)^2} N(\mu)$ ,  $K_3 = \frac{2}{3}\beta$ ,  $K_0 = 3K_3$ ,  $m^2 = -4N(\mu) \log \frac{T}{T_c}$ ,  $N(\mu) = \frac{\mu^2}{2\pi^2}$ ,  $g_s = \sqrt{\frac{24\pi^2 \lambda}{27 \log \mu/\Lambda}}$ ,  $T_c \sim \mu \exp \left( -\frac{3\pi^2}{\sqrt{2}g_s} \right)$ , where  $\mu$  is the chemical potential,  $\Lambda$  the QCD scale and  $T_c$  the critical temperature. The expectation value of  $\Phi$  can be directly found by minimizing the above potential (3) as:

$$\langle \Phi \rangle = \Delta_{\text{CFL}} \mathbf{1}_3, \quad \Delta_{\text{CFL}}^2 \equiv \frac{m^2}{8\beta}. \quad (4)$$

Masses of gauge bosons and scalars are given by the following:  $m_g^2 = g_s^2 \Delta_{\text{CFL}}^2 K_3 \lambda_3$ ,  $m_\zeta^2 = \frac{2m^2}{K_3}$ ,  $m_\chi^2 = \frac{4\lambda_2 \Delta_{\text{CFL}}^2}{K_3}$ ,  $m_\varphi^2 = 0$ , where  $\varphi$  is the massless Nambu-Goldstone boson related to the breaking of  $U(1)_B$  symmetry, and  $\zeta$  and  $\chi$  are respectively the trace and traceless part of  $\Phi$ . Static equations of motion can also be directly found from the Lagrangian above, and they read as:

$$\nabla_i F_{ij} = ig_s K_3 \lambda_3 \left[ \nabla_j \Phi \Phi^\dagger - \Phi (\nabla_j \Phi)^\dagger - \frac{1}{3} \text{Tr} (\nabla_j \Phi \Phi^\dagger - \Phi (\nabla_j \Phi)^\dagger) \right],$$

$$\nabla_j^2 \Phi = \frac{1}{K_3} [-m^2 + 2\beta \{ \Phi \Phi^\dagger + \text{Tr}(\Phi^\dagger \Phi) \}] \Phi. \quad (5)$$

<sup>1</sup> In this paper we are ignoring the first order time derivative term for simplicity since it makes the vortex dyonic.

### B. Non-Abelian vortex or color magnetic flux tube

Let us first briefly review a few primary features of the non-Abelian vortices in the CFL phase in the absence of the electromagnetic interaction. The transformation properties of the field  $\Phi$  can be written as

$$\Phi' = e^{i\theta_B} U_C \Phi U_F^{-1}, \quad e^{i\theta_B} \in U(1)_B, \quad U_C \in SU(3)_C, \quad U_F \in SU(3)_F. \quad (6)$$

There is a redundancy in the action of the discrete symmetries, and the actual symmetry group is given by

$$G = \frac{SU(3)_C \times SU(3)_F \times U(1)_B}{\mathbb{Z}_3 \times \mathbb{Z}_3}. \quad (7)$$

In the ground state  $\langle \Phi \rangle = \Delta_{\text{CFL}} \mathbf{1}_3$  with  $\Delta_{\text{CFL}} \equiv \sqrt{\frac{m^2}{8\beta}}$ , the full symmetry group  $G$  is spontaneously broken down to  $H \simeq \frac{SU(3)_{C+F}}{\mathbb{Z}_3}$ . The order parameter space is  $G/H \simeq \frac{SU(3) \times U(1)}{\mathbb{Z}_3} = U(3)$ . It can be easily noticed that  $\pi_1(G/H) = \mathbb{Z}$ . This non-zero fundamental group implies the existing vortices. Since the broken  $U(1)_B$  is a global symmetry, the vortices are global vortices or superfluid vortices [11]. The structure of these vortices can be understood by the orientation and winding of the configuration of the condensed scalar field  $\Phi$  in the far away from vortex core. We place a vortex along the  $z$ -direction and use the cylindrical coordinates in this paper. One can write down the Ansatz as

$$\Phi(r, \theta) = \Delta_{\text{CFL}} \begin{pmatrix} e^{i\theta} f_1(r) & 0 & 0 \\ 0 & f_2(r) & 0 \\ 0 & 0 & f_2(r) \end{pmatrix}, \quad A_i(r) = -\frac{1}{3g_s} \frac{\epsilon_{ij} x_j}{r^2} A(r) \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad i = \{1, 2\} \quad (8)$$

where  $f_1, f_2$  and  $A(r)$  are the profile functions.

The GL free energy can be written by inserting the Ansatz into Eq. (3) as

$$2\pi \int r dr \left[ \frac{2}{3g_s^2 \lambda_3 r^2} (\partial_r A)^2 + K_3 \Delta_{\text{CFL}}^2 \left\{ (\partial_r f_1)^2 + 2(\partial_r f_2)^2 + \frac{f_1^2}{9r^2} (3 - 2A)^2 + 2\frac{f_2^2}{9r^2} A^2 + \Delta_{\text{CFL}}^2 [f_1^2 - f_2^2]^2 + 2\Delta_{\text{CFL}}^2 [f_1^2 + 2f_2^2 - 3]^2 \right\} \right] \quad (9)$$

The form of the profiles  $f_1, f_2$  and  $A(r)$  can be calculated numerically with the boundary condition,

$$f_1(0) = 0, \quad \partial_r f_2(r)|_0 = 0, \quad A(0) = 0, \quad f_1(\infty) = f_2(\infty) = 1, \quad A(\infty) = 1. \quad (10)$$

The vortex configuration in Eq. (8) breaks the unbroken color-flavor diagonal  $SU(3)_{C+F}$  symmetry as

$$SU(3)_{C+F} \rightarrow SU(2) \times U(1) \quad (11)$$

showing the existence of degenerate solutions. This degeneracy is due to the existence of Nambu-Goldstone modes parameterizing a coset space  $\frac{SU(3)}{SU(2) \times U(1)} \simeq \mathbb{CP}^2$ . The low-energy excitation and interaction of these zero mode can be calculated by the effective  $\mathbb{CP}^2$  sigma model action [1]. Generic solutions on the  $\mathbb{CP}^2$  space can be found by just applying a global transformation by a reducing matrix[28],

$$U(\phi) = \frac{1}{\sqrt{X}} \begin{pmatrix} 1 & -\phi^\dagger \\ \phi & X^{\frac{1}{2}} Y^{-\frac{1}{2}} \end{pmatrix}, \quad X = 1 + \phi^\dagger \phi, \quad Y = \mathbf{1}_3 + \phi \phi^\dagger, \quad (12)$$

where  $\phi = \{\phi_1, \phi_2\}$  are inhomogeneous coordinates of the  $\mathbb{CP}^2$ . The vortex solution with a generic orientation takes the form,

$$\Phi(r, \theta) = \Delta_{\text{CFL}} U(\phi) \begin{pmatrix} e^{i\theta} f_1(r) & 0 & 0 \\ 0 & f_2(r) & 0 \\ 0 & 0 & f_2(r) \end{pmatrix} U^\dagger(\phi), \quad A_i(r) = -\frac{\epsilon_{ij} x_j}{3g_s r^2} A(r) U(\phi) \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} U^\dagger(\phi). \quad (13)$$

### III. THE CONSTRUCTION OF THE EFFECTIVE ACTION OF A NON-ABELIAN VORTEX

The effective action can be computed by prompting the moduli parameter  $\phi$  to fields fluctuating on the vortex world-sheet in the  $t$ - $z$  plane, which is known as the moduli approximation [25] (see also Refs. [26, 27]). So when one inserts the rotated solution in Eq. (13) into the free energy, the static energy part is separated out from the rests. The other terms which would be relevant for small fluctuations can be written as

$$\mathcal{L}_{\text{eff}} = \sum_{\alpha} c_{\alpha} \text{Tr} [F_{i\alpha} F_i^{\alpha} + \kappa_{\alpha} |\mathcal{D}^{\alpha} \Phi|^2] \quad (14)$$

where  $\alpha = \{0, 3\}$  is the world-sheet index,  $i = \{1, 2\}$ ,  $c_0 = \epsilon_3$ ,  $c_3 = \frac{1}{\lambda_3}$  and  $\kappa_{\alpha} = \frac{K_{\alpha}}{c_{\alpha}}$ . The raising and lowering of the index  $\alpha$  are done by the Minkowski signature  $(+, -)$  for  $\{0, 3\}$ . So the equations of motion for zero (the Gauss's law) and third component can be expressed as,

$$\mathcal{D}_i F^{i\alpha} = -ig_s \kappa_{\alpha} T^a \text{Tr} [\Phi^{\dagger} T^a \mathcal{D}^{\alpha} \Phi - (\mathcal{D}^{\alpha} \Phi)^{\dagger} T^a \Phi]. \quad (15)$$

These equations are generated due to the fluctuation of zero mode along the vortex. The Ansatz for generated gauge fields which solve the above Eq. (15) can be expressed as [32]

$$A_{\alpha} = \rho_{\alpha}(r, \theta) W_{\alpha} + \eta_{\alpha}(r, \theta) V_{\alpha}, \quad \alpha = 0, 3 \quad (16)$$

where  $\rho_{\alpha}$  and  $\eta_{\alpha}$  are profile functions which are to be determined by minimizing the action or by solving Eq. (15) and

$$W_{\alpha} = i\partial_{\alpha} \tilde{T} \tilde{T}, \quad V_{\alpha} = \partial_{\alpha} \tilde{T}, \quad \tilde{T} = UTU^{\dagger}, \quad T = \text{diag}(1, -1, -1), \quad (17)$$

where  $U(\phi)$  is defined in the last section Eq. (12). These satisfy the commutation relations:  $[W_{\alpha}, \tilde{T}] = 2iV_{\alpha}$ ,  $[V_{\alpha}, \tilde{T}] = -2iW_{\alpha}$ ,  $\text{Tr} W_{\alpha} W^{\alpha} = \text{Tr} V_{\alpha} V^{\alpha}$ ,  $\text{Tr} W_{\alpha} V^{\alpha} = 0$ . Here we can see that  $W_{\alpha}$  and  $V_{\alpha}$  are orthogonal to each other.  $W_{\alpha}$  and  $V_{\alpha}$  are also orthogonal to the direction of  $A_i$  defined in Eq. (13). We can see this if we expand all three matrices as

$$U(\phi) \frac{1}{3} \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} U^{\dagger}(\phi) = \frac{1}{6} \mathbf{1} + \frac{1}{2} \tilde{T}, \quad W_{\alpha} = i[\partial_{\alpha} U U^{\dagger} - \tilde{T} \partial_{\alpha} U U^{\dagger} \tilde{T}], \quad V_{\alpha} = -i[i\partial_{\alpha} U U^{\dagger}, \tilde{T}]. \quad (18)$$

$W_{\alpha}$  is the Delduc-Valent projection on  $\mathbb{CP}^2$  [28] and was used in singular gauge computation. Here we introduce  $V_{\alpha}$  as a new component in the Ansatz (16) to solve the Gauss's law. Getting  $A_{\alpha}$  orthogonal to  $A_i$  direction indicating the fluctuation of Nambu-Goldstone mode in the orthogonal direction of the background field. Which is true because Nambu-Goldstone bosons are generated by broken generators.

To compute the effective action we have to insert the Ansatz in Eq. (16) into the action in Eq. (14). Before we do so, let us compute the field strength of the gauge field and matter coupling separately. The first term of the field strength ( $F_{i\alpha} = \partial_i A_{\alpha} - \mathcal{D}_{\alpha} A_i$ ) can be written as

$$\partial_i A_{\alpha} = \left( \frac{x_i}{r} \partial_r \rho_{\alpha}(r, \theta) - \frac{\epsilon_{ij} x^j}{r^2} \partial_{\theta} \rho_{\alpha}(r, \theta) \right) W_{\alpha} + \left( \frac{x_i}{r} \partial_r \eta_{\alpha}(r, \theta) - \frac{\epsilon_{ij} x^j}{r^2} \partial_{\theta} \eta_{\alpha}(r, \theta) \right) V_{\alpha}. \quad (19)$$

The second term becomes

$$\mathcal{D}_{\alpha} A_i = \partial_{\alpha} A_i - ig_s [A_{\alpha}, A_i] = \frac{\epsilon_{ij} x^j}{r^2} A(r) \{ \eta_{\alpha}(r, \theta) W_{\alpha} - \sigma_{\alpha}(r, \theta) V_{\alpha} \} \quad (20)$$

where we have defined  $2g_s \sigma_{\alpha} = 1 + 2g_s \rho_{\alpha}$ . So we can insert the field strength

$$F_{i\alpha} = \partial_i A_{\alpha} - \mathcal{D}_{\alpha} A_i = \left( \frac{x_i}{r} \partial_r \rho_{\alpha} - \frac{\epsilon_{ij} x^j}{r^2} (\partial_{\theta} \rho_{\alpha} + A \eta_{\alpha}) \right) W_{\alpha} + \left( \frac{x_i}{r} \partial_r \eta_{\alpha} - \frac{\epsilon_{ij} x^j}{r^2} (\partial_{\theta} \eta_{\alpha} - A \sigma_{\alpha}) \right) V_{\alpha} \quad (21)$$

into the kinetic term of the gauge field in Eq. (14), to yield

$$\text{Tr} F_{i\alpha} F_i^{\alpha} = \left[ (\partial_r \rho_{\alpha})^2 + (\partial_r \eta_{\alpha})^2 + \frac{1}{r^2} (\partial_{\theta} \rho_{\alpha} + A \eta_{\alpha})^2 + \frac{1}{r^2} (\partial_{\theta} \eta_{\alpha} - A \sigma_{\alpha})^2 \right] \text{Tr} V_{\alpha} V^{\alpha}. \quad (22)$$

From the covariant derivative of the matter field  $\Phi$ ,

$$\begin{aligned} \mathcal{D}_{\alpha} \Phi &= \partial_{\alpha} \Phi - ig_s A_{\alpha} \Phi \\ &= \Delta_{\text{CFL}} \left[ \frac{f_1 e^{i\theta} - f_2}{2} (1 + g_s \rho_{\alpha}) - ig_s \eta_{\alpha} \frac{f_1 e^{i\theta} + f_2}{2} \right] V_{\alpha} - g_s \Delta_{\text{CFL}} \left( \eta_{\alpha} \frac{f_1 e^{i\theta} - f_2}{2} + i \rho_{\alpha} \frac{f_1 e^{i\theta} + f_2}{2} \right) W_{\alpha}, \end{aligned} \quad (23)$$

we compute the  $|\mathcal{D}_\alpha \Phi|^2$  as

$$\frac{|\mathcal{D}_\alpha \Phi|^2}{\Delta_{\text{CFL}}^2} = \frac{1}{4} \left[ (1 + 2g_s \rho_\alpha + 2g_s^2 \rho_\alpha^2)(f_1^2 + f_2^2) + 2g_s^2 \eta_\alpha^2 (f_1^2 + f_2^2) - (1 + 2g_s \rho_\alpha) 2f_1 f_2 \cos \theta - 4g_s f_1 f_2 \eta_\alpha \sin \theta \right] \text{Tr} V_\alpha V_\alpha. \quad (24)$$

By changing the variables from  $\rho_\alpha$  to  $\sigma_\alpha = \frac{1+2g_s \rho_\alpha}{2g_s}$ , the above Eq. (24) can be simplified as

$$\frac{4|\mathcal{D}_\alpha \Phi|^2}{\Delta_{\text{CFL}}^2} = \left[ \frac{1}{2}(f_1^2 + f_2^2) + 2g_s^2 (\sigma_\alpha^2 + \eta_\alpha^2) (f_1^2 + f_2^2) - 4g_s f_1 f_2 (\sigma_\alpha \cos \theta + \eta_\alpha \sin \theta) \right] \text{Tr} V_\alpha V_\alpha. \quad (25)$$

Let us define here a complex scalar field as

$$\Psi_\alpha(r, \theta) = \sigma_\alpha(r, \theta) + i\eta_\alpha(r, \theta) \quad (26)$$

and we can rewrite the action in terms of these fields. Eq. (22) becomes

$$\text{Tr} F_{i\alpha} F_i^\alpha = \sum_\alpha \left[ |\partial_r \Psi_\alpha|^2 + \frac{1}{r^2} |\mathcal{D}_\theta \Psi_\alpha|^2 \right] \text{Tr} W_\alpha W^\alpha \quad (27)$$

where  $\mathcal{D}_\theta = \partial_\theta - iA(r)$ . Eq. (25) can also be rewritten as

$$\frac{|\mathcal{D}_\alpha \Phi|^2}{\Delta_{\text{CFL}}^2} = \frac{1}{4} \left[ \frac{1}{2}(f_1^2 + f_2^2) + 2g_s^2 |\Psi_\alpha|^2 (f_1^2 + f_2^2) - 2g_s f_1 f_2 (\Psi_\alpha e^{-i\theta} + \Psi_\alpha^* e^{i\theta}) \right] \text{Tr} W_\alpha W^\alpha. \quad (28)$$

The effective Lagrangian in Eq. (14) can be written as

$$\mathcal{L}_{\text{eff}} = \sum_\alpha c_\alpha \int d^2x \text{Tr} [F_{i\alpha} F_i^\alpha + \kappa_\alpha |\mathcal{D}_\alpha \Phi|^2] = \sum_\alpha \mathcal{I}_\alpha \text{Tr} W_\alpha W^\alpha \quad (29)$$

where  $\mathcal{I}_\alpha$  are the coefficients of the  $\mathbb{CP}^2$  action, defined by

$$\mathcal{I}_\alpha = c_\alpha \int r dr d\theta \left[ |\partial_r \Psi_\alpha|^2 + \frac{1}{r^2} |\mathcal{D}_\theta \Psi_\alpha|^2 + \frac{\Delta_{\text{CFL}}^2 \kappa_\alpha}{4} \left\{ \frac{1}{2}(f_1^2 + f_2^2) + 2g_s^2 |\Psi_\alpha|^2 (f_1^2 + f_2^2) - 2g_s f_1 f_2 (\Psi_\alpha e^{-i\theta} + \Psi_\alpha^* e^{i\theta}) \right\} \right]. \quad (30)$$

The effective Lagrangian can be written explicitly as the form of the  $\mathbb{CP}^2$  model:

$$L_{\text{eff}} = \mathcal{I}_0 \{ \dot{\mathbf{n}}^\dagger \dot{\mathbf{n}} + (\mathbf{n}^\dagger \dot{\mathbf{n}})(\dot{\mathbf{n}}^\dagger \mathbf{n}) \} - \mathcal{I}_3 \{ \partial_z \mathbf{n}^\dagger \partial_z \mathbf{n} + (\mathbf{n}^\dagger \partial_z \mathbf{n})(\partial_z \mathbf{n}^\dagger \mathbf{n}) \} \quad (31)$$

where  $\mathbf{n}$  are the homogeneous coordinates of the  $\mathbb{CP}^2$  space, which can be written in terms of the inhomogeneous coordinates as

$$\mathbf{n} = \frac{1}{\sqrt{1 + \phi^\dagger \phi}} \begin{pmatrix} 1 \\ \phi \end{pmatrix}. \quad (32)$$

If we rescale the  $z$ -coordinate as

$$z' = \sqrt{\frac{\mathcal{I}_0}{\mathcal{I}_3}} z \quad (33)$$

then the effective Lagrangian becomes

$$L_{\text{eff}} = \mathcal{I}_0 \left[ \partial_\alpha \mathbf{n}^\dagger \partial^\alpha \mathbf{n} + (\mathbf{n}^\dagger \partial_\alpha \mathbf{n})(\partial^\alpha \mathbf{n}^\dagger \mathbf{n}) \right]. \quad (34)$$

By expanding the field  $\Psi_0$  in terms of partial waves as

$$\Psi_0 = \sum_m \Psi_m(r) e^{im\theta}, \quad (35)$$

$\mathcal{I}_0$  can be written in terms of the partial wave modes as

$$\mathcal{I}_0 = c_0 \sum_m \int r dr d\theta \left[ (\partial_r \Psi_m)^2 + \frac{(m - A(r))^2}{r^2} \Psi_m^2 + \frac{\Delta_{\text{CFL}}^2 \kappa_0}{4} \left\{ \frac{1}{2} (f_1^2 + f_2^2) + 2g_s^2 \Psi_m^2 (f_1^2 + f_2^2) - 4g_s f_1 f_2 \Psi_m \cos(m-1)\theta \right\} \right]. \quad (36)$$

It is easy to check that the last term of Eq. (36) vanishes after the theta integration unless  $m = 1$ . So we write  $\mathcal{I}_0$  as

$$\mathcal{I}_0 = \sum_m 2\pi c_0 \int r dr \left[ (\partial_r \Psi_m)^2 + \frac{(m - A(r))^2}{r^2} \Psi_m^2 + \frac{\Delta_{\text{CFL}}^2 \kappa_0}{4} \left\{ \frac{1}{2} (f_1^2 + f_2^2) + 2g_s^2 \Psi_m^2 (f_1^2 + f_2^2) - 4g_s f_1 f_2 \Psi_m \delta_{m1} \right\} \right]. \quad (37)$$

We minimize the above integral by solving the equations for the modes,

$$\frac{1}{r} \partial_r r \partial_r \Psi_1 - \frac{(1 - A(r))^2}{r^2} \Psi_1 = \frac{\Delta_{\text{CFL}}^2 \kappa_0 g_s}{2} [(f_1^2 + f_2^2) g_s \Psi_1 - f_1 f_2], \quad (38)$$

$$\frac{1}{r} \partial_r r \partial_r \Psi_m - \frac{(m - A(r))^2}{r^2} \Psi_m = \frac{\Delta_{\text{CFL}}^2 \kappa_0 g_s^2}{2} (f_1^2 + f_2^2) \Psi_m, \quad m \neq 1. \quad (39)$$

These equation can be also derived directly from the equations of motion in Eq. (15).

One should notice that Eq. (38) for  $m = 1$  was derived in the singular gauge in Ref. [1] (for  $K_0 = K_3$ ) but the rests  $m \neq 1$  were absent in the singular gauge. The solution of Eq. (38) is normalizable ( $m = 1$ ) and can be solved [1] numerically with the boundary condition  $\Psi_1(0) = 0$  and  $\Psi_1(\infty) = \frac{1}{2g_s}$ . Large distance behavior of the solution can be expressed as

$$\Psi_1 \rightarrow_{r \rightarrow \infty} \frac{1}{2g_s} - \frac{1}{\sqrt{\xi}} e^{-\xi} \quad (40)$$

and it is easy to show that the large distance value of  $\Psi_1 = \frac{1}{2g_s}$  transforms the Ansatz  $A_\alpha$  in Eq. (16) into a pure gauge form as

$$A_\alpha = \frac{i}{g_s} \mathbf{g}^\dagger \partial_\alpha \mathbf{g}, \quad \text{where} \quad \mathbf{g} = e^{-i\theta T_8^*}, \quad T_8^* = \frac{1}{3} U(\phi) \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} U^\dagger(\phi). \quad (41)$$

The coefficient  $\mathcal{I}_0(m = 1)$  can be written in terms of the solution of the equations of motion derived in the above:

$$\mathcal{I}(m = 1) = \frac{2\pi m_0^2 c_0}{8g_s^2} \int r dr [f_1^2 + f_2^2 - 4f_1 f_2 g_s \Psi_1], \quad (42)$$

where  $m_0^2 = g_s^2 \Delta_{\text{CFL}}^2 \kappa_0$ . The integrand vanishes at large distances, so the integral is finite.

All other modes ( $m \neq 1$ ) are non-normalizable divergent modes. In this work we do not solve these equations numerically. However, one can understand the behavior of the solutions once we analyze the asymptotic forms of the solutions. For the case  $m \neq 1$ , Eq. (39) can be expressed as

$$\frac{1}{r} \partial_r r \partial_r \Psi_m = \left[ \frac{(m - A(r))^2}{r^2} + \frac{\Delta_{\text{CFL}}^2 \kappa_0 g_s^2}{2} (f_1^2 + f_2^2) \right] \Psi_m. \quad (43)$$

The right hand side of the above equation is positive definite (by assuming  $\Psi_m$  is positive). So the solution cannot have a local maximum. At large distances the equation becomes the modified Bessel equation:

$$\xi^2 \Psi_m'' + \xi \Psi_m' - [(m-1)^2 + \xi^2] \Psi_m = 0 \quad (44)$$

where  $\xi = m_0 r$ . The solution is  $\Psi_m \sim \frac{1}{\sqrt{\xi}} e^{\pm \xi}$ .

At small distances near the origin ( $r = 0$ ) where  $f_1(r) = 0$ , Eq. (43) reduces to

$$\left( \partial_\xi^2 + \frac{1}{\xi} \partial_\xi \right) \Psi_m = \frac{m^2}{\xi^2} \Psi_m. \quad (45)$$

The solution is  $\Psi_m \sim \xi^{\pm m}$  for  $m \neq 0, 1$ . So  $\Psi_m(0) = 0$  for  $m \neq 0, 1$  and we may conclude that the solution diverges as  $\frac{1}{\sqrt{\xi}} e^\xi$  at large distances since it does not have any local maximum. For  $m = 0$ , near the origin the solution could be written as either  $\log \xi$  or  $C_0 + \xi^2 C_2$ , where  $C_0$  and  $C_2$  are constants and  $C_2$  is found to be positive. So the solution diverges at large distances even if we set the value as constant at the origin.



#### IV. SUMMARY AND DISCUSSION

In this paper, we have analyzed the orientational  $\mathbb{CP}^2$  zero modes of a single non-Abelian vortex in high density QCD. To do so, we have followed the standard procedure of zero mode analysis and have written the effective action, as was done in the singular gauge case. In order to solve the Gauss's law constraint in the regular gauge, we have generalized the Ansatz of the gauge field (used in the singular gauge) to  $A_\alpha = \rho(r, \theta)W_\alpha + \eta(r, \theta)V_\alpha$  by introducing a profile function  $\eta$  together with a matrix  $V_\alpha$  orthogonal to  $W_\alpha$ , both of which do not exist in the singular gauge. In the regular gauge, the two profile functions ( $\rho, \eta$ ) do not only depend on the radial coordinate  $r$  but also on the azimuthal angle  $\theta$ . These two profile functions can be combined to the real and imaginary components of a single complex profile function  $\Psi(r, \theta)$ . The insertion of  $A_\alpha$  together with vortex profile functions into the action gives the  $\mathbb{CP}^2$  effective action on the  $t$ - $z$  plane with a front coefficient depending on the complex profile function  $\Psi(r, \theta)$ . The front coefficient has been determined by inserting  $\Psi(r, \theta)$  after solving the equations of motion. We have expanded the complex profile function  $\Psi$  in the partial wave basis as  $\Psi = \sum_m \Psi_m(r)e^{im\theta}$  and have analyzed the asymptotic behaviors of all the modes. We have found that only one mode  $\Psi_1$  is normalizable, which is identical to what was found in the singular gauge analysis. We have shown that all the other modes diverge exponentially at large distances. Finally, we have concluded that our regular gauge analysis established the previously known result of the existence of normalizable zero mode derived in singular gauge and that the previously constructed effective  $\mathbb{CP}^2$  Lagrangian of the single vortex is correct.

Here let us discuss some points which may shed light on the interesting features of the regular gauge. In this analysis, we have introduced a complex profile function which does not depend only on  $r$  but also on the azimuthal angle  $\theta$ . This azimuthal angle dependence of a zero mode makes the system complicated. The  $\mathbb{CP}^2$  Nambu-Goldstone zero modes arise as a consequence of the unbroken color-flavor group  $SU(3)_{C+F}$  in the bulk, further broken as  $SU(3)_{C+F} \rightarrow U(1) \times SU(2)$  inside the vortex core. The system restores  $SU(3)_{C+F}$  symmetry at large distances from the vortex core, where the unbroken  $SU(3)_{C+F}$  group elements commute with the order-parameter. However, the presence of vortex makes the order-parameter position dependent at large distances. So the embedding of color-flavor diagonal group  $SU(3)_{C+F}$  inside the original symmetry group becomes space dependent. The generators of the  $SU(3)_{C+F}$  changes along a path around the vortex by the action of holonomy. The space dependence is true only for the few generators which belong to the  $\mathbb{CP}^2$  subspace of  $SU(3)$  and others remain unaffected. The azimuthal angle dependence of  $\mathbb{CP}^2$  generators actually makes the zero mode analysis complicated. So when we fluctuate the zero modes, the generated gauge field  $A_\alpha$  depends on the angle in a complicated way. In general, the generators may not return back to their own after a complete rotation along an encircled path around the vortex. In this case, it is said that there is the so-called obstruction, for which the profile functions in general diverges as  $r^c$  at large distances with a constant  $c$  [29–32]. In our case, after expanding our complex profile function in the partial wave basis, we have found that the profile functions corresponding to different partial wave modes diverge exponentially except for one normalizable mode ( $m = 1$ ). Therefore, as a byproduct of our analysis, we have shown that non-Abelian vortices in high density QCD do not suffer from any obstruction.

There is an alternative way to show the absence of normalizable modes other than the  $\mathbb{CP}^2$  modes and translational modes, that is, the index theorem. Fermionic zero modes around a single non-Abelian vortex [33, 34] were studied by the index theorem applied to the Bogoliubov-de Gennes equation [35]. The index theorem applied to bosonic modes should be studied in the framework of the GL theory.

#### ACKNOWLEDGEMENT

We would like to thank Mark Alford for bringing us an attention to a possible problem of the singular gauge in constructing the effective action. This work is supported by the Ministry of Education, Culture, Sports, Science (MEXT)-Supported Program for the Strategic Research Foundation at Private Universities “Topological Science” (Grant No. S1511006). C. C. acknowledges support as an International Research Fellow of the Japan Society for the Promotion of Science (JSPS). The work of M. N. is supported in part by JSPS Grant-in-Aid for Scientific Research (KAKENHI Grant No. 25400268), and by a Grant-in-Aid for Scientific Research on Innovative Areas “Topological Materials Science” (KAKENHI Grant No. 15H05855) and “Nuclear Matter in Neutron Stars Investigated by Experiments and Astronomical Observations” (KAKENHI Grant No. 15H00841) from the the Ministry of Education,

- 
- [1] M. Eto, E. Nakano and M. Nitta, “Effective world-sheet theory of color magnetic flux tubes in dense QCD,” *Phys. Rev. D* **80**, 125011 (2009) [arXiv:0908.4470 [hep-ph]].
  - [2] M. Eto, M. Nitta and N. Yamamoto, “Instabilities of Non-Abelian Vortices in Dense QCD,” *Phys. Rev. Lett.* **104**, 161601 (2010) [arXiv:0912.1352 [hep-ph]].
  - [3] M. G. Alford, K. Rajagopal and F. Wilczek, “QCD at finite baryon density: Nucleon droplets and color superconductivity,” *Phys. Lett. B* **422**, 247 (1998) [hep-ph/9711395].
  - [4] M. G. Alford, K. Rajagopal and F. Wilczek, “Color flavor locking and chiral symmetry breaking in high density QCD,” *Nucl. Phys. B* **537**, 443 (1999) [hep-ph/9804403].
  - [5] M. G. Alford, A. Schmitt, K. Rajagopal and T. Schäfer, “Color superconductivity in dense quark matter,” *Rev. Mod. Phys.* **80**, 1455 (2008) [arXiv:0709.4635 [hep-ph]].
  - [6] K. Rajagopal and F. Wilczek, “The Condensed matter physics of QCD,” In \*Shifman, M. (ed.): At the frontier of particle physics, vol. 3\* 2061-2151 (World Scientific) [hep-ph/0011333].
  - [7] K. Iida and G. Baym, “The Superfluid phases of quark matter: Ginzburg-Landau theory and color neutrality,” *Phys. Rev. D* **63**, 074018 (2001) [*Phys. Rev. D* **66**, 059903 (2002)] [hep-ph/0011229].
  - [8] K. Iida and G. Baym, “Superfluid phases of quark matter. 2: phenomenology and sum rules,” *Phys. Rev. D* **65**, 014022 (2002) [hep-ph/0108149].
  - [9] I. Giannakis and H. C. Ren, “The Ginzburg-Landau free energy functional of color superconductivity at weak coupling,” *Phys. Rev. D* **65**, 054017 (2002) [hep-ph/0108256].
  - [10] M. M. Forbes and A. R. Zhitnitsky, “Global strings in high density QCD,” *Phys. Rev. D* **65**, 085009 (2002) [hep-ph/0109173].
  - [11] A. P. Balachandran, S. Digal and T. Matsuura, “Semi-superfluid strings in high density QCD,” *Phys. Rev. D* **73**, 074009 (2006) [hep-ph/0509276].
  - [12] E. Nakano, M. Nitta and T. Matsuura, “Non-Abelian strings in high density QCD: Zero modes and interactions,” *Phys. Rev. D* **78**, 045002 (2008) [arXiv:0708.4096 [hep-ph]].
  - [13] E. Nakano, M. Nitta and T. Matsuura, “Non-Abelian Strings in Hot or Dense QCD,” *Prog. Theor. Phys. Suppl.* **174**, 254 (2008) [arXiv:0805.4539 [hep-ph]].
  - [14] M. Eto and M. Nitta, “Color Magnetic Flux Tubes in Dense QCD,” *Phys. Rev. D* **80**, 125007 (2009) [arXiv:0907.1278 [hep-ph]].
  - [15] M. Eto, Y. Hirono, M. Nitta and S. Yasui, “Vortices and Other Topological Solitons in Dense Quark Matter,” *PTEP* **2014**, no. 1, 012D01 (2014) [arXiv:1308.1535 [hep-ph]].
  - [16] M. G. Alford, S. K. Mallavarapu, T. Vachaspati and A. Windisch, “Stability of superfluid vortices in dense quark matter,” *Phys. Rev. C* **93**, no. 4, 045801 (2016) [arXiv:1601.04656 [nucl-th]].
  - [17] W. Vinci, M. Cipriani and M. Nitta, “Spontaneous Magnetization through Non-Abelian Vortex Formation in Rotating Dense Quark Matter,” *Phys. Rev. D* **86**, 085018 (2012) [arXiv:1206.3535 [hep-ph]].
  - [18] Y. Hirono and M. Nitta, “Anisotropic optical response of dense quark matter under rotation: Compact stars as cosmic polarizers,” *Phys. Rev. Lett.* **109**, 062501 (2012) [arXiv:1203.5059 [hep-ph]].
  - [19] C. Chatterjee and M. Nitta, “Aharonov-Bohm Phase in High Density Quark Matter,” *Phys. Rev. D* **93**, no. 6, 065050 (2016) [arXiv:1512.06603 [hep-ph]].
  - [20] A. Gorsky, M. Shifman and A. Yung, “Confined Magnetic Monopoles in Dense QCD,” *Phys. Rev. D* **83**, 085027 (2011) [arXiv:1101.1120 [hep-ph]].
  - [21] M. Eto, M. Nitta and N. Yamamoto, “Confined Monopoles Induced by Quantum Effects in Dense QCD,” *Phys. Rev. D* **83**, 085005 (2011) [arXiv:1101.2574 [hep-ph]].
  - [22] Y. Hirono, T. Kanazawa and M. Nitta, “Topological Interactions of Non-Abelian Vortices with Quasi-Particles in High Density QCD,” *Phys. Rev. D* **83**, 085018 (2011) [arXiv:1012.6042 [hep-ph]].
  - [23] M. Kobayashi, E. Nakano and M. Nitta, “Color Magnetism in Non-Abelian Vortex Matter,” *JHEP* **1406**, 130 (2014) [arXiv:1311.2399 [hep-ph]].
  - [24] H. Abuki, “BCS/BEC crossover in Quark Matter and Evolution of its Static and Dynamic properties: From the atomic unitary gas to color superconductivity,” *Nucl. Phys. A* **791**, 117 (2007) [hep-ph/0605081].
  - [25] N. S. Manton, “A Remark on the Scattering of BPS Monopoles,” *Phys. Lett.* **110B**, 54 (1982).
  - [26] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Manifestly supersymmetric effective Lagrangians on BPS solitons,” *Phys. Rev. D* **73**, 125008 (2006) [hep-th/0602289].
  - [27] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Solitons in the Higgs phase: The Moduli matrix approach,” *J. Phys. A* **39**, R315 (2006) [hep-th/0602170].
  - [28] F. Delduc and G. Valent, “Classical and Quantum Structure of the Compact Kahlerian  $\sigma$  Models,” *Nucl. Phys. B* **253**, 494 (1985).
  - [29] M. G. Alford, K. Benson, S. R. Coleman, J. March-Russell and F. Wilczek, “The Interactions and Excitations of Nonabelian Vortices,” *Phys. Rev. Lett.* **64**, 1632 (1990) Erratum: [*Phys. Rev. Lett.* **65**, 668 (1990)].



- [30] M. G. Alford, K. Benson, S. R. Coleman, J. March-Russell and F. Wilczek, “Zero modes of nonabelian vortices,” Nucl. Phys. B **349**, 414 (1991).
- [31] M. G. Alford, K. M. Lee, J. March-Russell and J. Preskill, “Quantum field theory of nonAbelian strings and vortices,” Nucl. Phys. B **384**, 251 (1992) [hep-th/9112038].
- [32] S. Bolognesi, C. Chatterjee, J. Evslin, K. Konishi, K. Ohashi and L. Seveso, “Geometry and Dynamics of a Coupled 4D-2D Quantum Field Theory,” JHEP **1601**, 075 (2016) [arXiv:1509.04061 [hep-th]].
- [33] S. Yasui, K. Itakura and M. Nitta, “Fermion structure of non-Abelian vortices in high density QCD,” Phys. Rev. D **81**, 105003 (2010) [arXiv:1001.3730 [hep-ph]].
- [34] C. Chatterjee, M. Cipriani and M. Nitta, “Coupling between Majorana fermions and Nambu-Goldstone bosons inside a non-Abelian vortex in dense QCD,” Phys. Rev. D **93**, no. 6, 065046 (2016) [arXiv:1602.01677 [hep-ph]].
- [35] T. Fujiwara, T. Fukui, M. Nitta and S. Yasui, “Index theorem and Majorana zero modes along a non-Abelian vortex in a color superconductor,” Phys. Rev. D **84**, 076002 (2011) [arXiv:1105.2115 [hep-ph]].